arXiv:1107.5889v3 [hep-ph] 24 Sep 2011

IN-MEDIUM MASS MODIFICATIONS OF THE D_0 AND B_0 MESONS WITH THE QCD SUM RULES

Zhi-Gang Wang[1] [1], Tao Huang[2] [2]
[1] Department of Physics, North China Electric Power University, Baoding 071003, P. R. China
[2] Institute of High Energy Physics and Theoretical Physics Center for Science Facilities, Chinese Academy of Sciences, Beijing 100049, P.R. China

Abstract

In this article, we calculate the in-medium mass modifications of the scalar mesons D_0 and B_0 using the QCD sum rules. In calculations, we observe that the D_0N and B_0N scattering lengths are about 1.1 fm and 4.1 fm respectively, the mass-shifts $\delta M_{D_0} = 69\,\mathrm{MeV}$ and $\delta M_{B_0} = 217\,\mathrm{MeV}$, and the D_0N and B_0N interactions are repulsive. The positive mass-shifts indicate that the decays of the higher charmonium states into the $D_0\bar{D}_0$ pair are suppressed.

PACS numbers: 12.38.Lg; 14.40.Lb; 14.40.Nd
Key Words: Nuclear matter, QCD sum rules

1 Introduction

The in-medium properties of hadrons play an important role in understanding the strong interactions, the heavy ion collisions and the nuclear astrophysics. There have been many experiments devoted to study the in-medium hadron properties. The upcoming FAIR (Facility for Antiproton and Ion Research) project at GSI provides the opportunity to extend the experimental studies into the charm sector. The CBM (compressed baryonic matter) collaboration intends to study the in-medium properties of the hadrons, including charmed mesons [1], while the $\bar{\mathrm{P}}$ANDA collaboration will focus on the charm spectroscopy, and mass and width modifications of the charmed hadrons in the nuclear matter [2]. Therefore, the in-medium properties of the charmed mesons become excellent subjects in recent years.

The suppression of the J/ψ production in relativistic heavy ion collisions (such as RHIC or LHC) may be one of the important signatures to identify the possible phase transition to the quark-gluon plasma [3]. Although the dissociation of the J/ψ in the quark-gluon plasma due to color screening reduces its production, the effects of hadron absorptions [4] and co-mover interactions [5] also play an important role, and one should be careful to make definitive conclusion. The in-medium modifications of the hadron properties can affect the productions of the open-charmed mesons and the J/ψ in the heavy ion collisions, as the higher charmonium states are considered as the major source of the J/ψ [6]. If the mass reductions of the D, $\bar{D}$ mesons are large enough, the excited charmonium states can decay to the $D\bar{D}$ pair instead of decaying to the lowest state J/ψ, the J/ψ itself can also decay to the $D\bar{D}$ pair at even higher nuclear density [7], although the excited charmonium states undergo mass modifications in the nuclear matter. The D-meson can also be produced through the reaction $\psi + N \to D + \bar{D}$ in the heavy ion collisions, on the other hand, the D-mesons couple strongly through the inelastic channels such as $DN \to \Lambda_c$, Σ_c [8].

The in-medium mass modifications of the $D, \bar{D}$ mesons have been studied with the QCD sum rules [8, 9], the quark-meson coupling model [10], the coupled-channel approach [11], etc. The works on the scalar D_0, $\bar{D}_0$ mesons are few [12, 13], it is interesting to study their mass modifications in the nuclear matter to see whether or not the decaying of the higher excited charmonium states to the $D_0\bar{D}_0$ pair is facilitated. Just like the D meson, the D_0 meson contains a charm-quark and a light quark. The existence of a light quark in the charmed mesons results in much difference between the in-medium mass modifications of the charmed-mesons and charmonium states. The former have large contributions from the light-quark condensates, while the latter are dominated by the gluon condensates [8, 9, 12, 14]. In this article, we study the mass modifications of the

---

[1] E-mail,wangzgyiti@yahoo.com.cn.
[2] Email: huangtao@ihep.ac.cn

$D_0, \bar{D}_0$ mesons in the nuclear matter using the QCD sum rules [15], furthermore, we study the corresponding mass modifications of the $B_0, \bar{B}_0$ mesons considering the heavy quark symmetry. In Ref.[12], Hilger and Kampfer study the in-medium mass modifications of the scalar $D_0, \bar{D}_0$ mesons and quantify the $D_0 - \bar{D}_0$ mass splitting using the QCD sum rules, the hadronic spectral density they choose differs from that of the present work. In Ref.[13], Tolos et al study the in-medium properties of the scalar mesons D_0, $D_{s0}(2317)$ and $X(3700)$ in the coupled-channel approach and reproduce them as dynamically-generated resonances.

The article is arranged as follows: we study the in-medium mass modifications of the scalar mesons D_0 and B_0 with the QCD sum rules in Sec.2; in Sec.3, we present the numerical results and discussions; and Sec.4 is reserved for our conclusions.

2 In-medium mass modifications of the D_0 and B_0 with QCD sum rules

We study the mass modification of the D_0 meson in nuclear matter with the two-point correlation function $\Pi(q)$. In the Fermi gas approximation for the nuclear matter, the $\Pi(q)$ can be divided into the vacuum part $\Pi_0(q)$ and the static one-nucleon part $\Pi_N(q)$, which is expected to be valid at relatively low nuclear density, and the $\Pi_N(q)$ can be approximated in the linear density of the nuclear matter [8, 16],

$$\Pi(q) = i\int d^4x\, e^{iq\cdot x}\langle T\left\{J(x)J^\dagger(0)\right\}\rangle_{\rho_N} = \Pi_0(q)+\Pi_N(q) \simeq \Pi_0(q)+\frac{\rho_N}{2M_N}T_N(q)\,, \qquad (1)$$

where the ρ_N denotes the density of the nuclear matter, and the forward scattering amplitude $T_N(q)$ is defined as

$$T_N(\omega,\boldsymbol{q}) = i\int d^4x e^{iq\cdot x}\langle N(p)|T\left\{J(x)J^\dagger(0)\right\}|N(p)\rangle\,, \qquad (2)$$

the $J(x)$ denotes the isospin averaged current,

$$J(x) = J^\dagger(x) = \frac{\bar{c}(x)q(x)+\bar{q}(x)c(x)}{2}, \qquad (3)$$

the q denotes the u or d quark, the $q^\mu = (\omega, \boldsymbol{q})$ is the four-momentum carried by the current $J(x)$, the $|N(p)\rangle$ denotes the isospin and spin averaged static nucleon state with the four-momentum $p = (M_N, 0)$, and $\langle N(\boldsymbol{p})|N(\boldsymbol{p}')\rangle = (2\pi)^3 2p_0\delta^3(\boldsymbol{p}-\boldsymbol{p}')$ [8]. The terms proportional to p_F^4, p_F^5, p_F^6, $\cdots$ can be neglected at the normal nuclear matter with the saturation density $\rho_N = \rho_0 = \frac{2p_F^3}{3\pi^2}$, as the Fermi momentum $p_F = 0.27\,\text{GeV}$ is a small quantity [16].

In the limit $\boldsymbol{q}\to\boldsymbol{0}$, the $T_N(\omega,\boldsymbol{q})$ can be related to the D_0N scattering T-matrix, $\mathcal{T}_{D_0N}(M_{D_0},0) = 8\pi(M_N+M_{D_0})a_{D_0}$, where the a_{D_0} is the D_0N scattering length. Near the pole position of the D_0-meson, the phenomenological spectral density $\rho(\omega,0)$ can be parameterized with three unknown parameters a, b and c [8],

$$\rho(\omega,0) = -\frac{f_{D_0}^2M_{D_0}^2}{\pi}\text{Im}\left[\frac{\mathcal{T}_{D_0N}(\omega,\boldsymbol{0})}{(\omega^2-M_{D_0}^2+i\varepsilon)^2}\right]+\cdots\,, \qquad (4)$$

$$= a\,\frac{d}{d\omega^2}\delta(\omega^2-M_{D_0}^2)+b\,\delta(\omega^2-M_{D_0}^2)+c\,\delta(\omega^2-s_0)\,, \qquad (5)$$

where the decay constant f_{D_0} is defined by $\langle 0|J(0)|D_0(k)+\bar{D}_0(k)\rangle = f_{D_0}M_{D_0}$, the terms denoted by $\cdots$ represent the continuum contributions. The first term denotes the double-pole term, and corresponds to the on-shell effect of the T-matrix, $a = -8\pi(M_N+M_{D_0})a_{D_0}f_{D_0}^2M_{D_0}^2$; the second term denotes the single-pole term, and corresponds to the off-shell effect of the T-matrix; and the third term denotes the continuum term or the remaining effects, where the s_0 is the continuum threshold. Then the mass-shift of the D_0-meson can be approximated as

$$\delta M_{D_0} = 2\pi\frac{M_N+M_{D_0}}{M_N M_{D_0}}\rho_N a_{D_0}\,. \tag{6}$$

In the low energy limit $\omega\to 0$, the $T_N(\omega,\mathbf{0})$ is equivalent to the Born term $T_N^{\rm Born}(\omega,\mathbf{0})$, i.e. $T_N(0)=T_N^{\rm Born}(0)$. We take into account the Born term at the phenomenological side,

$$T_N(\omega^2) = T_N^{\rm Born}(\omega^2)+\frac{a}{(M_{D_0}^2-\omega^2)^2}+\frac{b}{M_{D_0}^2-\omega^2}+\frac{c}{s_0-\omega^2}\,, \tag{7}$$

with the constraint

$$\frac{a}{M_{D_0}^4}+\frac{b}{M_{D_0}^2}+\frac{c}{s_0} = 0\,. \tag{8}$$

The contributions from the intermediate spin-$\frac{3}{2}$ charmed baryon states are zero in the soft-limit $q_\mu\to 0$ [17], and we only take into account the intermediate spin-$\frac{1}{2}$ charmed baryon states in calculating the Born term. The isospin states of the D_0-mesons have the D_0^0N and D_0^+N interactions,

$$\begin{aligned} D_0^0(c\bar{u})+p(uud)\ \text{or}\ n(udd) &\longrightarrow \Lambda_c^+,\Sigma_c^+(cud)\ \text{or}\ \Sigma_c^0(cdd)\,,\\ D_0^+(c\bar{d})+p(uud)\ \text{or}\ n(udd) &\longrightarrow \Sigma_c^{++}(cuu)\ \text{or}\ \Lambda_c^+,\Sigma_c^+(cud)\,, \end{aligned} \tag{9}$$

where $M_{\Lambda_c}=2.286\,\rm{GeV}$ and $M_{\Sigma_c}=2.454\,\rm{GeV}$ [18]. We can take $M_H\approx 2.4\,\rm{GeV}$ as the average value, where the H means either Λ_c^+, Σ_c^+, Σ_c^{++} or Σ_c^0. It is straightforward to calculate the Born term by writing down the Feynman diagram, the result is

$$T_N^{\rm Born}(\omega,\mathbf{0}) = \frac{2f_{D_0}^2M_{D_0}^2M_N(M_H-M_N)g_{D_0NH}^2}{\left[\omega^2-(M_H-M_N)^2\right]\left[\omega^2-M_{D_0}^2\right]^2}\,, \tag{10}$$

where the g_{D_0NH} denotes the strong coupling constants $g_{D_0N\Lambda_c}$ and $g_{D_0N\Sigma_c}$. On the other hand, there are no inelastic channels for the $\bar{D}_0^0N$ and D_0^-N interactions, i.e. $T_N^{\rm Born}(0)=0$.

We carry out the operator product expansion to the condensates up to dimension-5 at the large space-like region in the nuclear matter. Once analytical results at the level of quark-gluon degree's of freedom are obtained, then we set $\omega^2=q^2$, and take the quark-hadron duality and perform the Borel transform with respect to the variable $Q^2=-\omega^2$, finally obtain the following sum rule:

$$\begin{aligned} &a\left\{\frac{1}{M^2}e^{-\frac{M_{D_0}^2}{M^2}}-\frac{s_0}{M_{D_0}^4}e^{-\frac{s_0}{M^2}}\right\}+b\left\{e^{-\frac{M_{D_0}^2}{M^2}}-\frac{s_0}{M_{D_0}^2}e^{-\frac{s_0}{M^2}}\right\}+\frac{2f_{D_0}^2M_{D_0}^2M_N(M_H-M_N)g_{D_0NH}^2}{(M_H-M_N)^2-M_{D_0}^2}\\ &\left\{\left[\frac{1}{(M_H-M_N)^2-M_{D_0}^2}-\frac{1}{M^2}\right]e^{-\frac{M_{D_0}^2}{M^2}}-\frac{1}{(M_H-M_N)^2-M_{D_0}^2}e^{-\frac{(M_H-M_N)^2}{M^2}}\right\}=\left\{\frac{m_c\langle\bar{q}q\rangle_N}{2}\right.\\ &\left.-\langle q^\dagger iD_0q\rangle_N+\frac{m_c^2\langle q^\dagger iD_0q\rangle_N}{M^2}-\frac{2m_c\langle\bar{q}iD_0iD_0q\rangle_N}{M^2}+\frac{m_c^3\langle\bar{q}iD_0iD_0q\rangle_N}{M^4}\right\}e^{-\frac{m_c^2}{M^2}}\\ &+\frac{1}{16}\langle\frac{\alpha_sGG}{\pi}\rangle_N\int_0^1 dx\left(1+\frac{\widetilde{m}_c^2}{M^2}\right)e^{-\frac{\widetilde{m}_c^2}{M^2}}-\frac{1}{48M^4}\langle\frac{\alpha_sGG}{\pi}\rangle_N\int_0^1\frac{1-x}{x}\widetilde{m}_c^4e^{-\frac{\widetilde{m}_c^2}{M^2}}\,, \end{aligned} \tag{11}$$

where $\widetilde{m}_c^2=\frac{m_c^2}{x}$.

Differentiate above equation with respect to $\frac{1}{M^2}$, then eliminate the parameter b, we can obtain the sum rule for the parameter a. With the simple replacements $m_c\to m_b$, $D_0\to B_0$, $\Lambda_c\to\Lambda_b$ and $\Sigma_c\to\Sigma_b$, we can obtain the corresponding sum rule for the mass modification of the B_0 meson in the nuclear matter.

3 Numerical results and discussions

In calculations, we have assumed that the linear density approximation is valid at the low nuclear density, for a general condensate in the nuclear matter, $\langle\mathcal{O}\rangle_{\rho_N} = \langle 0|\mathcal{O}|0\rangle + \frac{\rho_N}{2M_N}\langle N|\mathcal{O}|N\rangle = \langle\mathcal{O}\rangle_0 + \frac{\rho_N}{2M_N}\langle\mathcal{O}\rangle_N$. The input parameters are taken as $\langle\bar{q}q\rangle_N = \frac{\sigma_N}{m_u+m_d}(2M_N)$, $\langle\frac{\alpha_s GG}{\pi}\rangle_N = -0.65\,\mathrm{GeV}(2M_N)$, $\langle q^\dagger iD_0 q\rangle_N = 0.18\,\mathrm{GeV}(2M_N)$, $\langle\bar{q}g_s\sigma Gq\rangle = 3.0\,\mathrm{GeV}^2(2M_N)$, $\langle\bar{q}iD_0iD_0q\rangle_N + \frac{1}{8}\langle\bar{q}g_s\sigma Gq\rangle_N = 0.3\,\mathrm{GeV}^2(2M_N)$, $m_u + m_d = 12\,\mathrm{MeV}$, $\sigma_N = 45\,\mathrm{MeV}$, $M_N = 0.94\,\mathrm{GeV}$, and $\rho_N = (0.11\,\mathrm{GeV})^3$ [19].

The hadronic parameters M_{D_0}, M_{B_0}, f_{D_0}, f_{B_0} are determined by the conventional two-point correlation functions $\Pi_0(q)$ using the QCD sum rules, $M_{D_0} = 2.355\,\mathrm{GeV}$, $M_{B_0} = 5.74\,\mathrm{GeV}$, $f_{D_0} = 0.334\,\mathrm{GeV}$, $f_{B_0} = 0.28\,\mathrm{GeV}$ with the threshold parameters $s_D^0 = 8.0\,\mathrm{GeV}^2$ and $s_B^0 = 39.0\,\mathrm{GeV}^2$, respectively. For the observed scalar meson $D_0(2400)$, the value $M_{D_0} = 2.355\,\mathrm{GeV}$ is consistent with the average of the experimental data $M_{D_0^0} = 2318\,\mathrm{MeV}$ and $M_{D_0^\pm} = 2403\,\mathrm{MeV}$ [18]. The uncertainties come from the hadronic parameters f_{D_0} and f_{B_0} can be approximated as $\frac{2\delta f_{D_0}}{f_{D_0}}$ and $\frac{2\delta f_{B_0}}{f_{B_0}}$, respectively.

In Fig.1, we plot the mass-shifts δM versus the Borel parameter M^2. From the figure, we can see that the values of the mass-shifts are rather stable with variations of the Borel parameter at the intervals $M^2 = (6.1-7.4)\,\mathrm{GeV}^2$ and $(33-39)\,\mathrm{GeV}^2$ in the charmed and bottom channels, respectively, the uncertainties originate from the Borel parameter M^2 are less than 1%. Furthermore, the exponential factor $e^{-\frac{s_0}{M^2}} < e^{-1}$ at those intervals, the continuum contributions are greatly suppressed. The main comtribtuions come from the terms $m_c\langle\bar{q}q\rangle_N$ and $m_b\langle\bar{q}q\rangle_N$, the operator product expansion is certainly convergent.

In the Borel windows, the mass-shifts $\delta M_{D_0} = 61\,\mathrm{MeV}$, $65\,\mathrm{MeV}$, $68\,\mathrm{MeV}$, $72\,\mathrm{MeV}$, $75\,\mathrm{MeV}$, $79\,\mathrm{MeV}$ and $\delta M_{B_0} = 213\,\mathrm{MeV}$, $215\,\mathrm{MeV}$, $217\,\mathrm{MeV}$, $219\,\mathrm{MeV}$, $221\,\mathrm{MeV}$, $223\,\mathrm{MeV}$ respectively at the values $g^2 = 0$, 20, 40, 60, 80, 100, where the g denotes the strong coupling constants $g_{D_0N\Lambda_c}$, $g_{D_0N\Sigma_c}$, $g_{B_0N\Lambda_b}$, $g_{B_0N\Sigma_b}$, the mass-shifts increase monotonously with increase of the squared strong coupling constants g^2.

The calculations based on the QCD sum rules indicate that the values of the strong coupling constants $g_{NN\sigma(600)} = 12 \pm 2$ and $g_{NNa_0(980)} = 11 \pm 2$ [20], which are of the same magnitude of the phenomenological value of the strong coupling constant $g_{NN\pi} = 13.5$. On the other hand, the value of the strong coupling constant $g_{DN\Lambda_c} = 6.74$ from the QCD sum rules is much smaller [21]. In this article, we take the approximation $g_{D_0N\Lambda_c} \approx g_{D_0N\Sigma_c} \approx g_{B_0N\Lambda_b} \approx g_{B_0N\Sigma_b} \approx 6.74$, and obtain the values $\delta M_{D_0} = 69\,\mathrm{MeV}$, $\delta M_{B_0} = 217\,\mathrm{MeV}$, $a_{D_0} = 1.1\,\mathrm{fm}$ and $a_{B_0} = 4.1\,\mathrm{fm}$.

The positive scattering lengths indicate that the D_0N and B_0N interactions are repulsive, it is difficult to form the D_0N and B_0N bound states. Which are in contrast to the DN and BN interactions, where the negative scattering lengths indicate that the DN and BN interactions are attractive, it is possible to form the DN and BN bound states.

Due to positive mass-shift δM_{D_0}, the decays of the high charmonium states to the $D_0\bar{D}_0$ pair obtain additional suppression in the phase space, and the decay into the lowest state J/ψ is preferred. While the negative mass-shift δM_D indicate that the high charmonium states decays to the $D\bar{D}$ pair are facilitated, and the production of the J/ψ is suppressed. The J/ψ production does not obtain additional suppression due to mass modification of the scalar meson $D_0(2400)$ in the nuclear matter.

In the present work and Ref.[8], the correlation functions $\Pi(q)$ are divided into the vacuum part and the static one-nucleon part, and the nuclear matter induced effects are extracted explicitly, while in Refs.[9, 12], the pole terms of the hadronic spectral densities are parameterized as $\frac{\mathrm{Im}\Pi(\omega,0)}{\pi} = F_+\delta(\omega - M_+) - F_-\delta(\omega + M_-)$, where $M_\pm = M \pm \Delta M$ and $F_\pm = F \pm \Delta F$, and the sum rules for the mass center M and the mass splitting ΔM are obtained. For the pseudoscalar $D, \bar{D}$ mesons, Hayashigaki obtains the mass-shift $\delta M_D = -50\,\mathrm{MeV}$ [8], while Hilger, Thomas and Kampfer obtain the mass-shift $\delta M_D = +45\,\mathrm{MeV}$ [9]. For scalar $D_0, \bar{D}_0$ mesons, the mass-shift



Figure 1: The mass-shifts δM versus the Borel parameter M^2, the (I) and (II) denote the D_0 and B_0 mesons respectively, the A, B, C, D, E and F correspond to the values $g^2 = 0$, 20, 40, 60, 80 and 100, respectively.

$\delta M_{D_0} = M - M_{D_0} < 0$ obtained by Hilger and Kampfer [12] differs from the result in the present work.

In those studies, irrespective of the parameterizations of the hadronic spectral densities, derivatives with respect to $1/M^2$ are used to obtain additional QCD sum rules so as to take special superpositions to delete the unknown parameters and extract the explicit expressions of the mass-shifts. The QCD sum rules from the derivatives are not necessarily work well, as the QCD sum rules are just a QCD model, the predictions can vary with the special assumptions. Furthermore, the isospin currents and isospin-averaged currents can also lead to differences in the hadronic spectral densities. All those predictions can be confronted with the experimental data in the future.

In Ref.[13], Tolos et al study the in-medium properties of the scalar mesons $D_0(2400)$, $D_{s0}(2317)$ and $X(3700)$ in the coupled-channel approach and reproduce them as dynamically-generated resonances, and observe that the $D_{s0}(2317)$ and $X(3700)$ enlarge their widths to the order of 100 and 200 MeV respectively at the normal nuclear matter, and the $D_0(2400)$ meson obtains an extra widening from the already large width, due to the D meson absorptions in the nuclear matter via the DN and DNN inelastic reactions. In the vacuum, the mass and width of the $D_0(2400)$ are $M_{D_0^0} = (2318 \pm 29)\,\text{MeV}$, $M_{D_0^\pm} = (2403 \pm 14 \pm 35)\,\text{MeV}$, $\Gamma_{D_0^0} = (267 \pm 40)\,\text{MeV}$, and $\Gamma_{D_0^\pm} = (283 \pm 24 \pm 34)\,\text{MeV}$, respectively [18]. For $D_0^0(2400)$, the spin-parity $J^P = 0^+$ assignment is favored, while the spin-parity J^P of the $D_0^\pm(2400)$ still needs confirmation [18]. The mass-shifts $\delta M_{D_0} = 69\,\text{MeV}$ and $\delta M_D = -50\,\text{MeV}$ obtained in the present work and Ref.[8] respectively favor the decays $D_0 \to D\pi$, we can expect that the in-medium width of the $D_0(2400)$ is larger, and our prediction is compatible with the observation of Ref.[13]. However, the large width disfavors the experimental observation of the relative small mass-shift.

4 Conclusion

In this article, we calculate the in-medium mass-shifts of the scalar mesons D_0 and B_0 using the QCD sum rules. At the low density of the nuclear matter, we can take the linear approximation, and extract the mass-shifts explicitly. Our numerical results indicate that the D_0N and B_0N scattering lengths are $a_{D_0} = 1.1\,\text{fm}$ and $a_{B_0} = 4.1\,\text{fm}$, respectively, the D_0N and B_0N interactions are repulsive, the mass-shifts are $\delta M_{D_0} = 69\,\text{MeV}$ and $\delta M_{B_0} = 217\,\text{MeV}$. The positive mass-shifts indicate that the J/ψ production does not obtain additional suppression due to the mass modification of the scalar meson $D_0(2400)$ in the nuclear matter.

Acknowledgments

This work is supported by National Natural Science Foundation, Grant Number 11075053, and the Fundamental Research Funds for the Central Universities.

References

[1] B. Friman et al, "The CBM Physics Book: Compressed Baryonic Matter in Laboratory Experiments", Springer Heidelberg.

[2] M. F. M. Lutz et al, arXiv:0903.3905.

[3] T. Matsui and H. Satz, Phys. Lett. **B178** (1986) 416; R. Vogt, Phys. Rept. **310** (1999) 197.

[4] B. Zhang et al, Phys. Rev. **C62** (2000) 054905; W. Cassing and E. L. Bratkovskaya, Nucl. Phys. **A623** (1997) 570; E. L. Bratkovskaya, W. Cassing and H. Stoecker, Phys. Rev. **C67** (2003) 054905.

[5] A. Capella and E. G. Ferreiro, Eur. Phys. J. **C42** (2005) 419; N. Armesto et al, Nucl. Phys. **A698** (2002) 583; S. Gavin and R. Vogt, Phys. Rev. Lett. **78** (1997) 1006.

[6] L. Antoniazzi et al, Phys. Rev. Lett. **70** (1993) 383; Y. Lemoigne et al, Phys. Lett. **B113** (1982) 509.

[7] Y. S. Golubeva et al, Eur. Phys. J. **A17** (2003) 275.

[8] A. Hayashigaki, Phys. Lett. **B487** (2000) 96.

[9] T. Hilger, R. Thomas and B. Kampfer, Phys. Rev. **C79** (2009) 025202.

[10] K. Tsushima et al, Phys. Rev. **C59** (1999) 2824; A. Sibirtsev, K. Tsushima and A. W. Thomas, Eur. Phys. J. **A6** (1999) 351.

[11] L. Tolos, J. Schaffner-Bielich and A. Mishra, Phys. Rev. **C70** (2004) 025203; L. Tolos, J. Schaffner-Bielich and H. Stoecker, Phys. Lett. **B635** (2006) 85; T. Mizutani and A. Ramos, Phys. Rev. **C74** (2006) 065201; L. Tolos, A. Ramos and T. Mizutani, Phys. Rev. **C77** (2008) 015207; L. Tolos, C. Garcia-Recio and J. Nieves, Phys. Rev. **C80** (2009) 065202.

[12] T. Hilger and B. Kampfer, Nucl. Phys. Proc. Suppl. **207-208** (2010) 277.

[13] L. Tolos, R. Molina, D. Gamermann and E. Oset, Nucl. Phys. **A827** (2009) 249c; L. Tolos, D. Gamermann, R. Molina, E. Oset and A. Ramos, arXiv:0905.1850.

[14] A. Hayashigaki, Prog. Theor. Phys. **101** (1999) 923; F. Klingl et al, Phys. Rev. Lett. **82** (1999) 3396; S. Kim and S. H. Lee, Nucl. Phys. **A679** (2001) 517; C. M. Ko and S. H. Lee, Phys. Rev. **C67** (2003) 038202; A. Kumar and A. Mishra, Phys. Rev. **C82** (2010) 045207.

[15] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. **B147** (1979) 385; L. J. Reinders, H. Rubinstein and S. Yazaki, Phys. Rept. **127** (1985) 1.

[16] E. G. Drukarev and E. M. Levin, Prog. Part. Nucl. Phys. **27** (1991) 77; E. G. Drukarev, M. G. Ryskin and V. A. Sadovnikova, Prog. Part. Nucl. Phys. **47** (2001) 73.

[17] Z. G. Wang, Eur. Phys. J. **C57** (2008) 711; Z. G. Wang, Eur. Phys. J. **C61** (2009) 299.

[18] K. Nakamura et al, J. Phys. **G37** (2010) 075021.

[19] T. D. Cohen et al, Prog. Part. Nucl. Phys. **35** (1995) 221; X. M. Jin et al, Phys. Rev. **C47** (1993) 2882; X. M. Jin et al, Phys. Rev. **C49** (1994) 464.

[20] T. M. Aliev and M. Savci, Phys. Rev. **D75** (2007) 045006.

[21] F. S. Navarra and M. Nielsen, Phys. Lett. **B443** (1998) 285.